

An Inter-Regional Comparison of Ozone Sensitivity to Reductions in Emissions in Central California

By

Su-Tzai Soong, Saffet Tanrikulu, Cuong Tran,
Yiqin Jia and Scott Beaver

Bay Area Air Quality Management District

AGU Fall Meeting, December 5-9, 2011

San Francisco, California

Motivation for this study

- Past attainment demonstrations have not accurately predicted future ozone amount
- One explanation may be that past modeling studies have underestimated the amount of ozone in Central California, especially in the San Joaquin Valley
- Recent improvements in models lead to better model performance
- The improved model performance is likely to give a more reliable attainment demonstration

Meteorological and Air Quality Models

Past model selections

- Meteorology model: MM5
- Air quality model: CAMx


Current model selections

- Meteorology model: WRF
- Air quality model: CMAQ

Episode selection: July 12-27, 2006

Model runs: base case and 20%, 40%, 60% NO_x and VOC reduction


Example of a past
MM5-CAMx
simulation



WRF-CMAQ results


Layer 1 O3 Model Domain

1-Hr O3 4PM PDT July 17, 2006




Layer 1 O3 Bay Area

1-Hr O3 4PM PDT July 17, 2006




Scatter Plot for SFB area July 17, 2006

Paired in cell and hour




Scatter Plot for SFB area July 17, 2006

Best in 3x3 cells and within 1 hour




Scatter Plot for SFB area July 17, 2006

Best in 5x5 cells and within 2 hours




Scatter Plot for SAC area July 17, 2006

Best in 5x5 cells and within 2 hours




Scatter Plot for SJV area July 17, 2006

Best in 5x5 cells and within 2 hours




RRF for Livermore

RRF for Livermore




RRF for Folsom

RRF for Folsom



RRF for Arvin

RRF for Arvin



Effect of NOx and VOC Reductions (%)

Livermore										
PPB					RRF					
VOC\NOx	0	20	40	60	VOC\NOx	0	20	40	60	
0	87.84	89.38	88.51	83.50	0	1.00	1.02	1.01	0.95	
20	84.01	85.76	85.46	81.38	20	0.96	0.98	0.97	0.93	
40	80.37	82.02	82.29	78.97	40	0.91	0.93	0.94	0.90	
60	76.92	78.17	78.83	76.22	60	0.88	0.89	0.90	0.87	
Folsom										
PPB					RRF					
VOC\NOx	0	20	40	60	VOC\NOx	0	20	40	60	
0	95.54	91.34	85.13	75.23	0	1.00	0.96	0.89	0.79	
20	93.66	89.85	84.00	74.65	20	0.98	0.94	0.88	0.78	
40	91.65	88.25	82.77	73.99	40	0.96	0.92	0.87	0.77	
60	89.56	86.46	81.40	73.19	60	0.94	0.90	0.85	0.77	
Arvin										
PPB					RRF					
VOC\NOx	0	20	40	60	VOC\NOx	0	20	40	60	
0	107.47	103.63	97.27	86.89	0	1.00	0.96	0.91	0.81	
20	103.63	100.37	94.75	85.34	20	0.96	0.93	0.88	0.79	
40	99.42	96.74	91.86	83.44	40	0.93	0.90	0.85	0.78	
60	94.89	92.72	88.55	81.14	60	0.88	0.86	0.82	0.75	

How to Achieve 8-hr Ozone NAAQS (% reductions)

	2006 DV	84 PPB			75 PPB		
		RRF	NO _x	VOC	RRF	NO _x	VOC
SFB	80				0.94	0	30
SAC	97	0.87	40	40	0.77	60	40
SJV	110	0.76	60	60	0.68	>60	>60

Additional Issues

Changes in the design values (ppb) and design value stations

	2004	2005	2006	2007	2008	2009	2010
SFB	84	78	80	77	81	78	80
SAC	102	97	97	98	102	100	102
SJV	116	113	110	107	108	105	104

Cool	102	97	95	96	98	93	89
Folsom	97	97	97	98	102	100	102